

# TILMICODEM 25

## ORAL SOLUTION FOR POULTRY



**NAME OF THE VETERINARY MEDICINAL PRODUCT:** TILMICODEM 25, as tilmicosin phosphate 250 mg/ml, oral solution for poultry.

**STATEMENT OF THE ACTIVE SUBSTANCE AND OTHER INGREDIENTS:** 1 ml solution contains: Tilmicosin phosphate: 250 mg (equivalent to 225 mg tilmicosin base), Excipients (propyl gallate, disodium edentate, orthophosphoric acid, purified water) ad. 1 ml.

### INDICATIONS

**In poultry:** for treatment of infections produced by susceptible bacteria such as: *Mycoplasma* spp, *Pasteurella multocida*, *Haemophilus paragalarum*, *Streptococcus* spp., anaerobs (*Clostridium* spp.) etc.

**TARGET SPECIES:** Poultry.

### DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

**In poultry:** the product is administered via drinking water in dosage of 15 - 20 mg tilmicosin/kg (equivalent to 17 – 22 mg tilmicosin phosphate/kg b.w.) corresponding to 0.07 – 0.09 ml Tilmicodem 25/kg b.w./day for 3 – 5 consecutive days. Based on the recommended doses and the number and weight of the animals to be treated, the exact daily amount of Tilmicodem 25 should be calculated according to the following formula:

$$\frac{\text{ml Tilmicodem 25 / kg body weight/day} \times \text{mean body weight of animals to be treated (kg)}}{\text{Mean daily water consumption (liters) / animal}} = \text{ml Tilmicodem 25 / liter of drinking water}$$

The uptake of medicated water is dependent on the clinical condition of animals. In order to obtain the correct dosage, the concentration in drinking water may have to be adjusted.

**ADVICE ON CORRECT ADMINISTRATION:** The product must be diluted before administration to the animals. During treatment, the animals should drink only medicated water. Medicated drinking water should be prepared fresh every 24 hours. To ensure a correct dosage, body weight should be determined as accurately as possible to avoid under dosing.

**WITHDRAWAL PERIOD:** Edible tissues of poultry: 12 days from cessation of oral medication. Not permitted for use in laying bird producing eggs for human consumption.

**SPECIAL PRECAUTIONS FOR STORAGE:** Keep away from reach and sight of children! Store in original packaging, at room temperature (15–25°C), protected from moisture and direct sunlight. Shelf life of the veterinary medicinal product as packaged for sale: 2 years. Shelf life after dilution in drinking water, according to directions: 24 hours. Shelf-life after first opening of package: 28 days. Close tight the recipient after use. Do not use after expiry date which is stated on the label.

**SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS:** For the environmental protection, medicines should not be disposed of via wastewater or household waste. Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

### DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED

**OTHER INFORMATION:** PE-HD flasks of 100 ml, 500 ml, and 1000 ml; PE-HD cans of 5 l, 10 l, and 20 l. Not all pack sizes may be marketed. For any information regarding this veterinary medicinal product, please contact the local representative of the marketing authorization holder.